



# Better2Know

100% CONFIDENTIAL ~ FAST RESULTS ~ WORLDWIDE

## Syphilis Fact Sheet

### What is Syphilis?

Syphilis is a sexually transmitted infection passed from person to person through direct bodily contact. It is commonly passed on through oral, vaginal or anal intercourse with someone infected with Syphilis or through contact with visible Syphilis sore, also called a Chancre.

### What are the symptoms (if any)?

Many people infected with Syphilis do not have any symptoms. They can remain at risk for late complications if they are not treated. The first symptoms of Syphilis may appear from 10-90 days (average 21 days) after infection.

There are three stages of Syphilis infection: Primary, Secondary and Latent. Primary is marked by a Chancre (sore) that, left untreated, progresses to the Secondary stage. The symptoms of this stage can include rashes, fever, swollen glands, sore throat, hair loss, headaches, and muscle aches. The latent stage progresses and, unknown to the carrier, may eventually damage internal organs, including the brain, nerves, eyes, heart, blood vessels, liver, bones, and joints.

### How is it treated?

Syphilis is curable and can be treated with antibiotics.

You should abstain from sexual intercourse until your treatment is finished, and ensure that your sexual partners are also treated to avoid re-infecting each other.

### What happens if you do not get treated?

If Syphilis is left untreated, there is an increased risk of getting other STIs, including HIV.




Image: CDC and Dr David Cox

*Syphilis: Treponema pallidum*

Those with untreated Syphilis can progress through all three stages of the disease.

In pregnant women, untreated Syphilis can cause miscarriages, premature births, stillbirths, or death of newborn babies. There is also a risk that the baby may have deformities, delays in development, or seizures, as well as other problems such as rashes, fever, swollen liver and spleen, anaemia, and jaundice. Rarely, the symptoms of Syphilis go unseen in infants so that they develop the symptoms of late-stage syphilis, including damage to their bones, teeth, eyes, ears, and brain.

### Better2Know Testing Information

If you have symptoms you may get tested for Syphilis at any time with a blemish screen. If you do not have any symptoms you should wait 9 days after any incident you are concerned about before getting tested for Syphilis.

Better2Know offers a choice of tests including an instant test which take only 20 minutes to run. Better2Know's standard and instant Syphilis tests use a blood sample. The blemish screen is a swab of a visible sore, spot or blemish for Syphilis.

---

Better2Know is the world's leading provider of private sexual health testing services. We will get you a clinic appointment or home testing kit quickly, and deliver fast and accurate results for your peace of mind. If you have any concerns about your sexual health please visit our website [www.better2know.com](http://www.better2know.com) or email us at [info@better2know.com](mailto:info@better2know.com) to book an appointment or purchase your test with us today.

© Better2Know Ltd 2017. Unit 8 Northgate Business Centre, White Lund Industrial Estate, Morecambe, LA3 3PA, United Kingdom

This leaflet is designed to provide you with basic information only. None of the information on this leaflet is to be used for the identification or diagnosis of any sexually transmitted disease or any other type of infection or condition.